
DRAFT

ENVIRONMENTAL ASSESSMENT

COOPERATIVE GYPSY MOTH ERADICATION PROJECT KING, JEFFERSON and LEWIS COUNTIES WASHINGTON

MARCH 1, 2004



Prepared by
Washington State Department of Agriculture
Plant Protection Division

In cooperation with
United States Department of Agriculture
Animal and Plant Health Inspection Service
Plant Protection and Quarantine



Published for the 2004 Gypsy Moth Eradication Season Washington State Department of Agriculture Plant Protection Division 3939 Cleveland Ave. SE Olympia WA 98501 1-800 443-6684

Inquiries regarding availability of this publication in alternative formats should be directed to the Laboratory Services Division at (360) 586-8489 or Telecommunications Device for the Deaf (360) 902-1996.

TABLE OF CONTENTS

I. PURPOSE AND NEED FOR ACTION	1
A. Decisions to be Made and Scope of Analysis	1
 Introduction	1 2
B. Proposed Action	3
C. Need For Action	3
Economic, Social, and Ecological Impacts Project Goals and Objectives	3 4
D. Authorizing Laws and/or Policies	5
State Authorizing Laws Federal Authorizing Laws Environmental Laws and Other Regulations	5
II. PUBLIC INVOLVEMENT AND ISSUES	5
A. Public Notification and Involvement B. Issues and Concerns	
III. AFFECTED ENVIRONMENT	8
A. 2004 Site Descriptions	8
B. Threatened, Endangered, and Sensitive Species	10
C. Other Environmental Consultation	11
IV. TREATMENT ALTERNATIVES	11
A. Treatment Alternatives	11
B. Preferred Treatment Alternative	11
C. Treatment Alternatives Not Selected	12

V. ENVIRONMENTAL CONSEQUENCES
A. Human Health and Safety13
1. Bacillus thuringiensis var. (kurstaki)132. Bond143. General Precautions15
B. Non-Target Organisms
1. Animals 15 2. Plants 16 3. Threatened, Endangered, and Sensitive Species 16
VI. MONITORING
VII. CUMULATIVE EFFECTS
VIII. SUMMARY
IX. LIST OF AGENCIES AND PERSONS CONSULTED 19
X. LIST OF PREPARERS20
XI. APPENDICES20
A. References
B. Treatment Site Maps
C. Alternatives Described in 1995 FEIS
D. Washington State Department of Health Recommendations
E. Product Labels & Material Safety Data Sheets
F. Standard Operating Procedures
G. Letters and Permits received through interagency consultation concerning

I. PURPOSE AND NEED FOR ACTION

A. Decisions to be Made and Scope of Analysis

1. Introduction

The Washington State Department of Agriculture (WSDA), in cooperation with the United States Department of Agriculture Animal and Plant Health Inspection Service (USDA-APHIS), is proposing an eradication program with the goal of eliminating an isolated infestation of the non-native gypsy moth, <u>Lymantria dispar</u> (Linnaeus), in King, Jefferson and Lewis Counties, Washington in the spring of 2004.

2. Environmental Analysis and Documentation

In 1995, the USDA Forest Service and APHIS prepared a final environmental impact statement, "Gypsy Moth Management in the United States: a cooperative approach", (hereinafter referred to as FEIS), which described and analyzed methods of gypsy moth control available for use in USDA cooperative programs. WSDA is proposing nothing that was not analyzed in the 1995 FEIS. Therefore, a new programmatic environmental impact statement will not be required.

This Environmental Assessment (EA) is "tiered" to the FEIS in accordance with the Council on Environmental Quality regulations for implementing the National Environmental Policy Act of 1969 (NEPA) (40 CFR 1502.20 and 40 CFR 1508.28). This EA provides the basic background information necessary for the site-specific analysis of the potential environmental effects of WSDA's proposed 2004 Cooperative Gypsy Moth Eradication Project. The FEIS and this site-specific EA jointly constitute the environmental analysis and documentation required under NEPA.

Copies of the FEIS and the EA are available for review at:

Washington State Library Point Plaza East, Bldg. 1 6880 Capitol Blvd. S Tumwater, WA 98501

and

USDA, APHIS, PPQ APHIS Library, 1st floor 4700 River Road Riverdale, MD 20737

and

USDA, APHIS, PPQ 22000 Marine View Drive S., Suite 201 Des Moines, WA 98198

Additional environmental analysis and documentation has been prepared to satisfy Washington State requirements under Chapter 43.21 (c) of the Revised Code of Washington (State Environmental Policy Act or SEPA), and Chapter 197-11 of the Washington Administrative Code (SEPA rules).

Copies of the SEPA documentation are available for review at:

Washington State Library Point Plaza East, Bldg. 1 6880 Capitol Blvd. S Tumwater, WA 98501

3. History and Scope of Project

Since its accidental release in the United States in 1869, the European strain of gypsy moth has spread throughout New England and areas to the north, south and west. It has become established in all or parts of 19 states, the District of Columbia, and parts of Canada. It continues to spread to uninfested areas. The gypsy moth has caused dramatic economic, social, and ecological impacts throughout the infested area (USDA, 1995, vol. II, chapter 1, p. 4).

The European strain of the gypsy moth has been found every year in Washington State since 1974 with the exceptions of 1976 and 1977. The European gypsy moth is usually introduced to Washington State by people visiting or relocating from the infested area of eastern North America. For more than 25 years, WSDA has successfully detected and eradicated new introductions of gypsy moth.

In 1991, the Asian strain of the gypsy moth was found for the first time in Oregon, Washington, and in British Columbia, Canada. Eradication projects conducted in 1992 successfully eliminated the insect from those areas. WSDA has detected and treated introductions of the Asian strain of the gypsy moth in 1991-92, 1994-95, 1995-96, 1996-97, 1997-98 and 1999-2000. These eradication projects have been successful. The Asian strain poses a far greater risk of rapid spread than the European. Unlike females of the European strain, females of the Asian strain may fly and deposit an egg mass miles from where they feed as caterpillars. The Asian strain also poses a greater risk of damage because it feeds on a greater variety of plants (USDA, 1995, vol. II, chapter 1, p. 4).

In 2004, WSDA is proposing to treat three sites that have isolated populations of the European strain of the gypsy moth. One is in the city of Bellevue in King County, the second is in the Port Ludlow area of Jefferson County and the third is in the Mayfield area of Lewis County.

For more information on how the different strains/populations of the gypsy moth are to be treated please see USDA, 1995, vol. II, chapter 1, pp. 9-11.

4. Decisions to be Made

There are three significant decisions, which must be made as a part of evaluating a gypsy moth control action.

The first decision to be made is whether to propose a gypsy moth control project (the absence of a control project is a no-action alternative). The second decision to be made is whether or not tiering this environmental assessment to the USDA 1995 FEIS is appropriate. The third decision to be made is whether to proceed with the preferred alternative as described in the FEIS.

B. Proposed Action

Strategies described in the FEIS depend upon the infestation status of the area: generally infested, transition, or uninfested. The three strategies of suppression, eradication, and slow the spread -- or their absence – are included in the six alternatives described in the FEIS. The sixth alternative is the preferred alternative presented in the FEIS. The sixth alternative is comprised of all three strategies.

Based on the infestation status of "no established population" Washington State's strategy in 2004 will be eradication.

For a more detailed description of the alternatives described in the FEIS, please refer to an excerpt from the FEIS in Appendix C of this EA.

Treatments available for eradication projects include: (the biological insecticides) Bacillus thuringiensis var. kurstaki (B.t.k.) and the gypsy moth nucleopolyhedrosis virus (Gypchek); a chemical insecticide (diflubenzuron); and treatments employing mass trapping, mating disruption, and sterile insect release techniques. A detailed description of these treatments is available in Appendix A of the FEIS.

C. Need For Action

1. Economic, Social, and Ecological Impacts

In order to avoid undesirable economic, social, and ecological impacts to residents, communities and businesses in Washington State, WSDA in cooperation with USDA APHIS, proposes to eradicate three isolated infestations of European gypsy moth. One is in the city of **Bellevue** in King County, the second is in the **Port Ludiow** area of Jefferson County and the third is in the **Mayfield** area of Lewis County.

Trapping (utilizing pheromone-baited traps) and visual inspections for alternate life stages such as egg masses have detected gypsy moth infestations in the aforementioned areas. The gypsy moth is able to survive and reproduce in Washington State, as evidenced by numerous past isolated infestations. The current infestations, if left unchecked, could spread across large areas.

Trees in forests and orchards, and residential and municipal shade trees and landscape plantings would be damaged and killed. Recreational and aesthetic values associated with trees and forested land would be diminished (USDA, 1995, vol. II, chapter 2, p. 29). Species composition of the vegetation on forested land could change, affecting the quantity and variety of food available for wildlife (USDA, 1995, vol. II, chapter 2, p. 23).

Water quality could be adversely affected in a number of ways including: 1) increased siltation from rapid runoff of rainfall from defoliated areas; 2) increases in water temperature as it flows through areas made shadeless; and 3) nutrient overloading from the deposition of large quantities of caterpillar droppings (USDA, 1995, vol. II, chapter 2, pp. 24-25).

The pesticide load in the environment would likely increase in quantity, variety, and net detrimental environmental impact as home and business owners respond to ever-increasing numbers of gypsy moth caterpillars, the damage they cause, and the nuisance they represent (USDA, 1995, vol. II, chapter 4, p. 76).

Human health effects associated with the presence of large numbers of gypsy moth caterpillars have been reported, including rashes and welts typical of allergic reactions, and respiratory complaints. These effects have been attributed to the irritating nature of the bristles found on the caterpillars. In some instances the reactions have been severe enough to require medical attention (USDA, 1995, vol. III, chapter 3, pp. 2-3), (Allen et, al., 1991), (Tuthill, et al., 1984), (Aber, et al., 1982), (Beaucher and Farnham, 1982), (Shama, et al., 1982).

Agricultural, horticultural and forestry enterprises are dependent upon markets beyond the borders of Washington State. Washington must be able to comply with the plant pest and disease regulations of the Federal government, other states, and international markets. The establishment and spread of the gypsy moth in Washington State would result in the imposition of quarantines (USDA, 1995, vol. II, chapter 2, p. 29). The levels of production and value of plant products would be adversely affected.

2. Project Goals and Objectives

The WSDA, in cooperation with USDA-APHIS and other appropriate Federal, State and local agencies, proposes to take action to eradicate three isolated infestations of European gypsy moth. One site is in the city of Bellevue in King County, the second is in the Port Ludlow area of Jefferson County and the third is in the Mayfield area of Lewis County. The action will be designed to give the project the best chance for

achieving the goal of eradicating the gypsy moth infestations while minimizing risks to human health as well as minimizing detrimental environmental consequences. This action will be taken in order to prevent the establishment and spread of this pest insect and thereby avoid the adverse economic, social, and ecological effects associated with large-scale gypsy moth infestations.

D. Authorizing Laws and/or Policies

1. State Authorizing Laws

WSDA has authority under Chapter 17.24 of the Revised Code of Washington, Insect Pests and Plant Diseases, to eradicate or control insect pests that may endanger the agricultural and horticultural industries in the state of Washington.

2. Federal Authorizing Laws

The USDA-APHIS has broad discretionary authority to prevent the establishment or spread of plant pests. See 1995 FEIS, volume 2, chapter 1, pages 8 and 9, "Statutory Authorities", for more information.

3. Environmental Laws and Other Regulations

Many environmental laws, authorities and Executive Orders of the President influence how actions to manage pests, including the gypsy moth, are implemented at the site-specific level. Such laws include the National Environmental Policy Act; the Washington State Environmental Policy Act; the Federal Insecticide, Fungicide, and Rodenticide Act; the Clean Water Act and the Endangered Species Act. See 1995 FEIS, volume 2, chapter 1, pages 8 and 9, "Statutory Authorities", for more information.

II. PUBLIC INVOLVEMENT AND ISSUES

A. Public Notification and Involvement

In the **early fall of 2003** Washington State Department of Agriculture (WSDA) employees made personal contact with some residents and businessmen at Port Ludlow, Bellevue, and Mayfield. This contact was made as the employees searched for egg masses and other evidence of gypsy moth activity at the three sites where multiple catches of moths had been made earlier in the summer.

On December 15, 2004 letters were sent to stakeholders in Port Ludlow, Bellevue, and Mayfield.

The letters said the following: 1) A gypsy moth infestation has been located in the community. 2) WSDA is proposing to eradicate the infestation the site with the biological insecticide *Bacillus thuringiensis* var. *kurstaki (Btk)*. 3) WSDA will soon begin

a public information campaign to inform local residents and community leaders of the infestation and proposed treatment.

The stakeholders to whom letters were sent included state legislators, county commissioners, county council members, mayors, and city council members.

On December 18, 2003 WSDA published a news release announcing that they were proposing action in the spring of 2004 to prevent the European gypsy moth from becoming established at an 18-acre site in Port Ludlow, an 11-acre site in Bellevue, and 7.5-acre site in Mayfield.

The news release emphasized the following: 1) Before a decision is made on the WSDA proposal, WSDA will prepare a State Environmental Policy Act (SEPA) checklist and National Environmental Policy Act (NEPA) environmental assessment for public review and comment, and consult other state and federal agencies on the proposal. 2) Residents whose properties were located in the proposed treatment zones will receive written information on the infestation and proposed treatment. 3) Citizens are encouraged to call the WSDA toll-free hotline (1-800-443-6684) or visit the WSDA Web site (www.agr.wa.gov, and click on "gypsy moth") if they have any questions about the infestation or proposed eradication. 4) Opens houses will be held in the tree communities in February so residents can review display boards, pick up written information, view a videotape, and ask questions of entomologists regarding the proposal.

In late December and early January numerous news articles appeared in Port Ludlow, Bellevue, and Mayfield area newspapers as a result of the news release being dispatched.

On January 22, 2004 letters with three enclosures were sent to residents in Port Ludlow and Mayfield, and to businesses in Bellevue (there are no residents in the proposed Bellevue treatment zone).

The letters: 1) Contained details of the infestation. 2) Informed residents and businesses that WSDA was proposing to eradicate the problem by treating the site with a biological insecticide, *Bacillus thuringiensis* var. *kurstaki (Btk)* 3) Provided the time and place when an open house would be held in their community so they could gather additional information on the infestation and proposal. Enclosed to the letter were a gypsy moth fact sheet, map of the proposed treatment site, and a copy of the news release sent earlier to the media.

On February 3, 2004 the media and stakeholders were provided electronically with an update of WSDA's proposal to eradicate gypsy moth infestations in Port Ludlow, Bellevue, and Mayfield.

Stakeholders were informed of the following: 1) Letters had been sent January 22, 2004 to residents and businesses containing details of the infestation and proposed

treatment and inviting them to an open houses to be held in their community. 2) A gypsy moth open houses would be held in the community at a date, time, and place listed in the letter. 3) Two environmental documents detailing the impact of the proposed treatment on the environment would be posted to WSDA's Web site (www.agr.wa.gov) by early March. The two documents would also be made available in hard copy at a local library (the address and telephone number of the library were provided). 4) Three key points being stressed by WSDA in conversations with residents and community leaders were: a) The gypsy moth was first detected in Washington in 1974 but a permanent population has never been established. b) WSDA has conducted 73 eradication projects since 1979 and all have been successful. c) The biological insecticide proposed for use [Btk] has a proven safety record.

On February 9, 2004 an open house was held in Port Ludlow at the North Bay Beach Club.

On February 11, 2004 an open house was held in Bellevue in the cafeteria at Sammamish High School.

On February 17, 2004 an open house was held in Mayfield in the library at Mossyrock High School.

On February 18, 2004 letters were sent to residents in Port Ludlow, and Mayfield, and to businesses in Bellevue requesting permission to come on their property beginning in late April or early May and apply three treatments of a biological insecticide, 7 to 14 days apart. Permission could be granted one of three ways: 1) Calling the WSDA toll-free hotline (1-800-443-6684) and giving permission verbally. 2) Completing a form enclosed in the letter and mailing it to WSDA in self-addressed envelope also enclosed in the letter. 3) Faxing the completed form to WSDA at (360) 586-8509.

B. Issues and Concerns

Concerns were raised about the proposed treatments, their effects on human health and on non-target organisms. Those issues raised are addressed in this EA and in the FEIS to which this EA is "tiered".

III. AFFECTED ENVIRONMENT

A. 2004 SITE DESCRIPTIONS (see Appendix B for maps)

Bellevue (Bellevue South, WA 7.5X15 minute quadrangle, S28, 33 T25N R5E)

- King County, Washington
- Approximately 11 acres
- Zoning: Commercial Office
- Approximately 15 properties in the proposed treatment area.
- Proposed Boundaries: The boundaries of the proposed site are: on the west, just west of 119th Ave NE; on the east, just east of 120th Ave NE; on the north, approximately 425ft north of NE 8th St; and on the south, approximately 200ft south of NE 8th St.
- Vegetation

The proposed treatment area is primarily comprised of deciduous trees. Canopy coverage is less than 5%, tree height is variable with trees up to 60 feet.

Critical/Sensitive Areas:

None within the proposed treatment area however, Lake Sturtevant/Bellevue is 150 feet to the north Sturtevant creek is 250 feet to the west. And a 40% Slope is 150 feet to the east

- Catch History
 - 1 European Gypsy Moth was caught in the area during the 2001 summer survey. 6 European Gypsy Moth were caught in the area during the 2002 summer survey. 21 European Gypsy Moths were caught in the area during the 2003 summer survey.
- Alternate Life Stages
 6 European Gypsy Moth egg masses and numerous pupal cases were found in this area in the late summer of 2003.

Port Ludlow (Port Ludlow, WA 7.5 minute quadrangle, S17 T28N R1E)

- Jefferson County, Washington
- Approximately 18 acres
- Zoning: MPR Single Family

- Approximately 32 properties in the proposed treatment area.
- Proposed Boundaries: The western boundary is Osprey Ridge Dr.; the northern boundary is Goldfinch Ln.; the eastern boundary lies just to the west of Rainer Ln.; and the southern boundary is just to the north of Hummingbird Ct.
- Vegetation

The proposed treatment area is a mix of deciduous and coniferous trees growing in and around residential properties and a small canyon. Canopy coverage is approximately 50%, tree height is variable with deciduous trees in excess of 100 feet.

Critical/Sensitive Areas:

Type 4 Stream (seasonal) Slight Landslide Hazard

Catch History

14 European Gypsy Moths were caught in this area during the 2003 summer survey.

• Alternate Life Stages

3 European Gypsy Moth egg masses, numerous pupal cases, cast skins and a dead female Gypsy Moth were found in this area in the late summer of 2003.

Mayfield (Mayfield Lake, WA 7.5 minute quadrangle, S35 T13N R2E)

- Lewis County, Washington
- Approximately 7.5 acres
- Zoning: RDD 1-5 (Rural Development District 1 house/5 acres)
- Approximately 2 properties in the proposed treatment area.
- Proposed Boundaries: The boundaries of the proposed site are: to the east Flynn Rd, the remaining boundaries extend in a half circle with a radius of 450 feet to the north, south and west of a residential property to the west of Flynn Rd. The northern boundary is just to the south of Ciannigan Hill Rd.
- Vegetation

The proposed treatment area is primarily deciduous trees growing in and around a residential property and a farm. Canopy coverage is less than 10%, tree height is variable with deciduous trees up to 70 feet.

Critical/Sensitive Areas:

None within the proposed treatment area however, An undeliniated Wetland is 300 feet to the southwest

- Catch History
 8 European Gypsy Moths were caught in this area during the 2003 summer survey.
- Alternate Life Stages
 3 European Gypsy Moth pupal cases and one cast skins were found in this area in the fall of 2003.

B. Threatened, Endangered, and Sensitive Species

As required by the Endangered Species Act of 1973, the USDA is taking part in section 7 consultation with both the United States Fish and Wildlife Service (USFWS) and the National Oceanic and Atmospheric Administration Fisheries Service (NOAA Fisheries). In addition the WSDA has consulted with the Washington State Department of Fish and Wildlife (WDFW) and the Washington State Department of Natural Resources (DNR). These agencies provided maps or other data intended to aide in the identification of habitats of concern and the presence of listed, proposed, candidate, threatened or endangered species. See Appendix G.

The information provided by WDFW Priority Habitats and Species Program did not identify any threatened or endangered species on the **Bellevue** site, however, a number of raptor nesting sites were listed as occurring in the area. All listed raptor nesting sites are well over one half mile from the proposed site. Also, listed were records of bald eagle breeding occurrence well over one mile from the proposed site. WDFW also listed the presence of priority anadromous fish and priority resident fish in the area. The priority anadromous fish listed include fall chinook, coho salmon and sockeye salmon. The priority resident fish listed include resident cutthroat and rainbow trout.

The information provided by WDFW Priority Habitats and Species Program did not identify any threatened or endangered species on the **Port Ludlow** site, however, osprey, great blue heron and purple martin nesting sites were listed as occurring in the area. All listed nesting sites were well over one quarter mile from the proposed site. Also, listed was a record of a bald eagle breeding occurrence well over one half mile from the proposed site. WDFW also listed the presence of priority anadromous fish and priority resident fish in the area. The priority anadromous fish listed include chum salmon, coho salmon and winter steelhead. The only priority resident fish listed was resident cutthroat.

The information provided by WDFW Priority Habitats and Species Program did not identify any threatened or endangered species on the **Mayfield** site, however, osprey and bald eagle nesting sites were listed as occurring in the area. All listed nesting sites were well over one half mile from the proposed site. Also, listed was a record of a federal species of concern van dyke's salamander well over one mile from the proposed site. In addition elk and bald eagle were listed as concentrated while

harlequin ducks, osprey and bald eagle were listed as a breeding occurrence. WDFW also listed the presence of priority resident fish in the area. The priority resident fish listed include resident cutthroat and rainbow trout.

A retrieval of information from the WDFW butterfly database did not name any threatened, endangered or sensitive species on or near this site. See Appendix G.

The DNR Washington Natural Heritage Program reviewed their Natural Heritage database. The DNR found no records for rare plants or high quality ecosystems in the vicinity of this project. See Appendix G.

C. Other Environmental Consultation

The USDA is taking part in Endangered Species Act Section 7 consultation with both the United States Department of Interior Fish and Wildlife Sevice (USFWS) and the National Oceanic and Atmospheric Administration Fisheries Service (NOAA Fisheries).

The WSDA is taking part in a water quality consultation with the Washigton State Department of Ecology (DOE).

IV. TREATMENT ALTERNATIVES

A. Treatment Alternatives

WSDA is proposing to conduct an Integrated Pest Management (IPM) program to eradicate gypsy moth in Washington State. Integrated Pest Management involves selecting those options and techniques that give the best chance of meeting the project goal of eradication. The FEIS contains a range of alternatives from which WSDA has selected an IPM strategy. The treatment alternatives detailed in the FEIS include:

- 1. Bacillus thuringiensis var. kurstaki (B.t.k.)
- 2. Diflubenzuron
- 3. Gypchek
- 4. Mass trapping
- 5. Mating disruption
- 6. Sterile release

B. <u>Preferred Treatment Alternative</u>

The WSDA/USDA-APHIS gypsy moth eradication project IPM strategy proposed for 2004 includes the use of the biological insecticide (B.t.k.) Foray 48B (EPA Reg. No.

73049-46). This insecticide may be mixed with the spreader-sticker Bond. Treatments will also include visual inspections for and removal of egg masses when found, and be followed up by delimitation trapping. This IPM strategy will give the project the best chance to achieve the goal of eradicating the gypsy moth infestations while minimizing risks to human health and minimizing detrimental environmental consequences. Details of the proposed ground applications follow:

Ground applications:

The proposed action would involve three applications of <u>Bacillus thuringiensis</u> var. <u>kurstaki</u> (B.t.k.) with ground-based equipment at a rate of approximately 24 Billion International Units (BIU) per acre to foliage within the designated treatment areas. The applications would be made 3-14 days apart and would occur during the period between April 15 and June 30, 2004. Provisions would be made for a fourth ground application of B.t.k. if substantial rainfall occurs too soon following the completion of an application. Exact timing of the applications would be dependent on development of gypsy moth larvae and/or foliage as determined by WSDA.

A spreader-sticker (Bond) may be utilized, as an adjuvant at a rate of up to 16 ounces per 100 gallons of tank mix. Mixing the formulation with adjuvants for gypsy moth eradication projects has been common practice (USDA, 1995, vol. II, A-4).

All ground applications would be conducted in accordance with all applicable federal, state, and local laws and regulations, and would adhere to the Standard Operating Procedures developed by WSDA for this project. See Appendix F.

Follow up:

A follow up trapping program employing pheromone-baited traps in the summer of 2004 would contribute to the success of the eradication project by removing males from any residual population, delimiting the location of any residual populations of Gypsy moths, and aiding in the evaluation of the project.

In the event of multiple moth catches in a treatment area, visual inspections for alternate life stages (egg masses etc.) would be performed in the fall of 2004 to aid in determine if re-treatment actions should be considered.

C. Treatment Alternatives Not Selected

The remaining treatment alternatives available for this proposed eradication project, as outlined in the FEIS, were not selected due to lack of availability, unproven efficacy, or environmental/biological concerns (USDA, 1995, vol. II, pp. A3-10).

V. <u>ENVIRONMENTAL CONSEQUENCES</u>

A. Human Health and Safety

1. Bacillus thuringiensis var. (kurstaki) (B.t.k.)

The use of B.t.k. for the eradication of isolated gypsy moth infestations is expected to have no adverse impact on human health or the environment. Various strains of Bacillus thuringiensis (B.t.) are a naturally occurring bacterial component of soils worldwide. Modern aqueous formulations of B.t.k. used in gypsy moth control projects contain no organic solvents and have an excellent safety record associated with their use in gypsy moth suppression and eradication projects. An exemption from the requirement of a tolerance has been established for residues of B.t.k. in or on all raw agricultural commodities. This exemption stipulates that manufacturers of B.t.k. test each lot for pathogenicity and vertebrate toxicity. See Appendix E for a Sample Label and MSDS.

A detailed discussion of the human health effects of B.t.k. may be found in the 1995 FEIS vol. II, chapter 4, pp. 13-17, and in vol. III, chapter 4.

Due to advances in scientific knowledge, the law requires that pesticides which were first registered before November 1, 1984 be reregistered to ensure that they meet today's more stringent standards. In March of 1998 the United States Environmental Protection Agency came out with a Reregistration Eligibility Decision (EPA, 1998) in which they concluded:

Based on the reviews of the generic data for the active ingredient *Bacillus thuringiensis*, the Agency has sufficient information on the health effects of *Bacillus thuringiensis* and on its potential for causing adverse effects in fish and wildlife and the environment. The Agency has determined that *Bacillus thuringiensis* products, manufactured, labeled and used as specified in this Reregistration Eligibility Decision, will not pose unreasonable risks or adverse effects to humans or the environment. Therefore, the Agency concludes that products containing *Bacillus thuringiensis* for all uses are eligible for reregistration (EPA, 1998).

In the spring of 1999, Foray 48B was applied by aircraft to 52 square miles of Southern Vancouver Island to combat an infestation of European gypsy moth. Approximately 80,000 residents lived in the spray zones. The Capital Health Region coordinated a human health study of possible short-term health effects. The resulting report (Capital Health Region, 1999) concluded:

The results of this project did not show a relationship between aerial spraying of Foray 48B and short-term human health effects. Although some people self-reported health problems that they attributed to the spray program, the research

and surveillance methods used in this project did not detect any change in health status that could be linked to the spray program. Our results showed that many of the health complaints people reported during the spray were as common in people before the spray as they were shortly after the spray. This conclusion is consistent with those of previous studies of the possible health effects of Btk-based pesticide spray programs.

Exposure to B.t.k. spray resulting from its use as proposed in this gypsy moth eradication project is unlikely to cause significant human health effects. However, it is good practice to minimize exposure to any insecticide. One of the conclusions reached in the Oregon study by Green, et al.(1990), was that, "the level of risk for B.t.k. and other existing or future microbial pesticides in immunocompromised hosts deserves further study."

In addressing the issue of exposure to immunocompromised individuals and the general public to the treatments proposed in Spring 2004, the following recommendations were made by the Washington State Department of Health in February, 2004 (Appendix D).

The Washington State Department of Health (DOH) recommends that people in the area to be sprayed minimize exposure by doing the following:

- 1. Stay indoors for at least 30 minutes after the spraying to allow droplets to settle.
- 2. Wait until the spray has dried before letting skin touch the treated leaves and bushes.
- 3. Wash skin with soap and water if you come in contact with the spray.
- 4. People in the sprayed area can sign up with the Department of Agriculture (800-443-6684) to be notified the day before spraying. (WSDOH, 2004, see Appendix D)

2. Bond

Bond may be used as an adjuvant with the insecticide utilized in this proposed eradication program. Bond is a non-ionic spreader-sticker which acts as an adjuvant when mixed with insecticides. Bond is not an eye or primary skin irritant per the Federal Hazardous Substances Labeling Act. In the unlikely event that overexposure were to occur, local irritation might be possible, especially in sensitive individuals. Systemic toxic effects are unlikely. See Appendix E for a Sample Label and MSDS.

3. General Precautions

The WSDA will take the following additional steps to assist the public in avoiding or reducing exposure to the spray material:

- 1. The Pesticide Sensitive Individuals database, maintained by the Pesticide Management Division of the WSDA, will be checked for people living in or near the proposed treatment area who require advance notification.
- 2. The WSDA will offer a toll-free telephone line with information regarding scheduled treatment days.
- 3. The WSDA will provide notification calls the day before scheduled applications to any resident in the proposed treatment area requesting them.
- 4. During ground treatments WSDA on-site spray block monitors will notify residents before the actual application to their property.
- 5. During ground treatments WSDA on-site spray block monitors will notify bicyclists, joggers and other pedestrians that they are approaching the treatment area.
- 6. Information will be provided to residents of the treatment area about how to avoid or reduce exposure to the spray material.

B. Non-Target Organisms

1. Animals

Bacillus thuringiensis var. (kurstaki) (B.t.k.)

A detailed discussion of the ecological effects of B.t.k. on non-target organisms may be found in the 1995 FEIS vol. II, chapter 4, pp. 52-55, and in vol. IV, chapter 5, pp. 5-10.

As used in gypsy moth eradication projects, B.t.k. has not been shown to adversely affect fish, birds, mammals, or most non-target insects, including honey bees (USDA, 1995, vol. II, chapter 4, pp. 54-55). It is expected that B.t.k. may kill other lepidopteran larvae (leaf-eating caterpillars) if they are present in project areas when treatments occur. In turn, animals dependent on caterpillars as food theoretically may be affected. However, reductions in native caterpillar populations are expected to be temporary due to the brief residual effectiveness of B.t.k. deposits on foliage (4 to 10 days), the high reproductive capacity of most lepidoptera, and recolonization from adjacent untreated areas (USDA, 1995, vol. II, chapter 4, pp. 54-55). The small size of the proposed treatment areas should aid in the recolonization process.

A study conducted in Oregon in connection with gypsy moth control programs in 1986 and 1987 found reduced numbers of caterpillars immediately following B.t.k. treatments and reduced species diversity. This study also found that recovery in numbers of non-target caterpillars began the same season, but that recovery of species diversity lagged behind (Miller, 1990).

One study has shown that B.t.k. could interfere with the biological control of the noxious weed tansy ragwort by cinnabar moth larvae if applied to areas where the weed occurs when late-instar larvae are active (James, et al., 1993). However, an intentionally introduced species of flea beetle has more impact as the primary biological control agent on tansy ragwort (L.C. Burrill, et al. 1994). It is not anticipated that this proposed project would have any adverse impact on flea beetle populations.

Two studies examined the indirect effect of B.t.k. on the reproductive success of insectivorous birds through a possible reduction in food supply. The studies reported no significant differences between treated and untreated areas in numbers of eggs hatched or in nestling growth and development. When caterpillars weren't available, the birds switched to other available prey (Gaddis, 1987), (Gaddis and Corkran, 1986).

There is no evidence of significant adverse impacts of B.t.k. on aquatic organisms. In a study conducted on a benthic stream community there was no evidence that addition of B.t.k. to stream mesocosms created adverse effects for these communities even at greater than 100 times expected exposure rates (Richardson and Perrin, 1994).

2. Plants

Bacillus thuringiensis var. (kurstaki) (B.t.k.)

B.t.k. is non-toxic to plants. B.t.k. is sensitive to meteorological effects once it has been applied to plant surfaces. B.t.k. is readily removed from plant surfaces by rain and is rapidly degraded by sunlight (USDA, 1995, vol. IV, chapter 7, pp. 15). The use of Bond will help slow the removal and degradation of B.t.k. by both rain and sunlight.

Changes in soil productivity and fertility due to B.t.k. are not likely. B.t.k. persists for a relatively short time, B.t. is known to occur naturally in soils worldwide, and applications of insecticides containing B.t. do not appear to increase levels of B.t. in soil (USDA, 1995, vol. I, p. 19). For more information about the fate of B.t.k. in the soil refer to 1995 FEIS, vol. 4, chapter 7, p. 16.

3. Threatened, Endangered, and Sensitive Species

No threatened, endangered, or sensitive species are known to be in or near the proposed treatment sites. In reference to the species listed in the Affected Environment section of this EA all occur well outside of the proposed treatment sites. Therefore, it is not anticipated that the proposed use of B.t.k. would adversely effect these named species.

VI. MONITORING

During the treatment operation, a WSDA-designated monitor will observe all mixing and application of the spray material to ensure compliance with all federal, state, and local laws and regulations and adherence to the Standard Operating Procedures. See Appendix F.

The treatment sites will be intensively monitored in the summer of 2004 and 2005 using pheromone-baited traps to determine the effectiveness of the treatment, assist in the eradication and delimit any residual populations of gypsy moths. This monitoring may indicate a need for further action.

VII. CUMULATIVE EFFECTS

No cumulative effects due to the proposed action are anticipated.

VIII. SUMMARY

This EA has analyzed the potential environmental effects of the proposed WSDA and USDA APHIS treatment program. This analysis was based on the 1995 USDA FEIS entitled, "Gypsy Moth Management in the United States: a cooperative approach" and the preferred alternative strategy proposed by the Washington State Department of Agriculture and USDA-APHIS for eradicating Gypsy moths at three sites in Washington State. The WSDA/USDA-APHIS gypsy moth eradication project strategy proposed for 2004 includes the use of the biological insecticide (B.t.k.) and the spreader-sticker Bond, followed up by trapping, visual inspections and removal of egg masses where appropriate. It is believed that this IPM strategy will give the project the best chance of achieving the goal of eradicating the gypsy moth infestations while minimizing risks to human health and the environment.

To summarize:

- A. B.t.k. used as described in this Environmental Assessment presents minimal risk of significant impact on human health.
- B. It is not anticipated that any non-target animal or plant populations would be adversely affected due to the limited size of the treatment areas. Any detrimental effects on susceptible non-target organisms would be transient and these populations would recover as individuals from nearby untreated areas re-colonized the treatment areas.
- C. No threatened, endangered, or sensitive species would be adversely affected by this eradication project.

- D. No detrimental effects on vegetation, water, or soil are known or anticipated due to this eradication project.
- E. No cumulative effects are known or anticipated.

IX. LIST OF AGENCIES AND PERSONS CONSULTED/NOTIFIED

United States Department of Agriculture, Animal and Plant Health Inspection Service, Dr. Clinton Campbell, on content and style of EA.

Washington State Department of Health, Barbara Morrissey, for review of the proposed treatment with regard to human health concerns.

Washington State Department of Natural Resources, Natural Heritage Program, Ms. Sandy Swope Moody, for review of the proposed treatment area for the presence of sensitive species or habitats.

Washington State Department of Natural Resources, Mr. Dave Cole, for determination and advice with regard to forest practices at the Jefferson Co. site.

Washington State Department of Fish and Wildlife, Ms. Lori Guggenmos, for review of the proposed treatment area for the presence of sensitive species or habitats.

Washington State Department of Fish and Wildlife, Ms. Ann Potter, for review of the proposed treatment area for the presence of sensitive lepidopteran species.

Washington State Department of Fish and Wildlife, Mr. Steve Manlow, Lewis Co. regional habitat program manager.

Washington State Department of Fish and Wildlife, Mr. Rich Costelo, King Co. regional habitat program manager.

Washington State Department of Fish and Wildlife, Mr. Steve Kalinowski, Jefferson Co. regional habitat program manager.

X. <u>LIST OF PREPARERS</u>

C. H. Phillips, J. W. Townsend & J. L. Cena Entomologists Washington State Department of Agriculture 3939 Cleveland Ave. SE Olympia, WA 98501 1-800-443-6684

XI. APPENDICES

- A. References
- B. Treatment Site Maps
- C. Alternatives Described in 1995 FEIS
- D. Washington State Department of Health Recommendations
- E. Product Labels & Material Safety Data Sheets
- F. Standard Operating Procedures
- G. Letters and Permits received through interagency consultation concerning threatened, endangered, and sensitive species and habitats

APPENDIX A

REFERENCES

Aber, R., DeMelfi, T., Gill, T., Healey, B., Oswell, N., Ruhig, W., Speziale, H., and Witte, E. J. 1982. Rash illness Associated with Gypsy Moth Caterpillars--Pennsylvania. Morbidity and Mortality Weekly Report. 31:13, pp. 169-70.

Allen, V.T.; Gredmiller, O.; Tyler, W.B. 1991. Gypsy Moth Caterpillar Dermatitis Revisited. <u>Journal of the American Academy of Dermatology</u>. 24(6/1): 979-981.

Anderson, J.F.; Furniss, W.E. 1983. Epidemic of Urticaria Associated with First-instar Larvae of the Gypsy Moth (Lepidoptera: Lymantriidae). <u>Journal of Medical Entomology</u>. 20(2): 146-150.

Beaucher, W. N., and Farnham, J.E. 1982. Gypsy-Moth-Caterpillar Dermatitis. <u>The New England Journal of Medicine</u>. 306:21, pp. 1301-02.

Burrill, L.C., Callihan, R.H., Parker, R., Coombs, E., Radtke, H. 1994. Tansy Ragwort Senecio jocobaea L. A Pacific Northwest Extension Publication, Oregon, Idaho, Washington.

Capital Health Region Office of the Medical Health Officer Director of Research. 1999. Human Health Surveillance During the Aerial Spraying for Control of North American Gypsy Moth on Southern Vancouver Island, British Columbia, 1999. December 31, 1999, p 34.

Corbett, E.S. 1992. Gypsy Moth Defoliation Impacts on Water Quality and Quality. U.S. Department of Agriculture Interagency Gypsy Moth Research Forum. University Park PA: U.S. Department of Agriculture, Forest Service, Northeastern Forest Experiment Station, 310 Forest Resources Laboratory. p. 31.

Duphar, B.V. 1987. Fact Sheet on Dimilin, Report No. C.303.0503.

Eisler R. 1992. Diflubenzuron Hazards to Fish, Wildlife, and Invertebrates: A Synoptic Review. U.S. Dept. of Interior, Fish and Wildlife Service, Biological Report 4: Contaminant Hazard Reviews Report 25, June 1992.

Farlow, J.E. 1976. Dimilin the Aquatic Fauna of a Louisiana Coastal Marsh. Ph.D. Thesis, Louisiana State University and Agricultural and Mechanical College, Baton Rouge. 144 pp.

- Gaddis, P.K., and Corkran, C.C. 1986. Secondary Effects of B.t. Spray on Avian Predators: the Reproductive Success of Chestnut-backed Chickadees.
- Gaddis, P.K. 1987. Secondary Effects of B.t. Spray on Avian Predators: the Reproductive Success of Chickadees.
- Gartrell, M. 1981. Diflubenzuron. U.S. Food and Drug Administration, Bureau of Foods, HFF-420. 8 pp.
- Green, M., Heumann, M., Sokolow, R., Foster, L.R., Bryant, R., and Skeels, M. 1990. Public Health Implications of the Microbial Pesticide <u>Bacillus thuringiensis</u>: An Epidemiological Study, Oregon, 1985-86. <u>American Journal of Public Health</u>. 80:7, pp. 848-52.
- Ivie, G. W. 1978. Fate of Diflubenzuron in Cattle and Sheep. <u>Journal of Agriculture and Food Chemistry</u> .26:90-94.
- James, R.R., Miller, J.C., and Lighthart, B. 1993. <u>Bacillus thuringiensis</u> var. <u>kurstaki</u> Affects a Beneficial Insect, the Cinnabar Moth (Lepidoptera: Arctiidae). <u>Journal of Economic Entomology</u>. 86:2, pp.334-39.
- Kubena, L.F. 1981. The Influence of Diflubenzuron on Several Weight Characteristics in Growing Male Broiler and Layer Chickens. <u>Poultry Science</u>. 60:1175-1182.
- Kubena, L.F. 1982. The Influence of Diflubenzuron on Several Reproductive Characteristics in Male and Female Layer-Breed Chickens. <u>Poultry Science</u>. 61:268-271.
- Martinat, P.J., Christman, V., Cooper, R.J., Dodge, K.M., Whitmore, R.C., Booth, G., and Seidel, G. 1987. Environmental Fate of Dimilin 25-W in a Central Appalachian Forest. Bulletin of Environmental Contamination and Toxicology. 39:142-149.
- Miller, J.C. 1990. Field Assessment of the Effects of a Microbial Pest Control Agent on Nontarget Lepidoptera. <u>American Entomologist</u>. 37, pp. 135-39.
- Miller, R.W., Corley, C., Oehler, D.D., and Pickens, L.G. 1976. Feeding TH 6040 to Cattle: Residues in Tissues and Milk and Breakdown in Manure. <u>Journal of Agricultural and Food Chemistry</u>. 24:687-688.
- Muzzarelli, R. 1986. Chitin Synthesis Inhibitors: Effects on Insects and on Nontarget Organisms. CRC Critical Reviews in Environmental Control. 16:141-146.
- National Gypsy Moth Management Group, Inc. 1991. A Rash of Gypsy Moths: Allergic Reactions to Caterpillars a Serious Problem. Newsletter (Spring); p 3.

Noble, M.A., Riben, P.D., and Cook, G.J. 1992. Microbiological and Epidemiological Surveillance Programme to Monitor the Health Effects of Foray 48B BTK Spray.

Novo Nordisk, Commonly Asked Questions Concerning Foray, The Environment and the Safety of Foray. Foray Information.

Perlman, F. 1965. Arthropods in Respiratory Tract Allergy: Their Relationship to Allergens in House Dust. Acta Allergologica (Copenhagen) 21: 241-253.

Petsonk E. 1994. National Institute for Occupational Safety and Health, Montgomery, WV. [Telephone conversation with Patrick R. Dunkin]. 7 March.

Richardson, John S.; Perrin, Chris J. 1994. Effects of the Bacterial Insecticide Bacillus thuringiensis var. kurstaki (B.t.k.) on a Stream Benthic Community. Canadian Journal of Aquatic Sciences. 51.

Richmond, M.L., Henny, C.J., Floyd, R.L., Mannan, R.W., Finch, D.M., and DeWeese, L.R. 1979. Effects of Sevin-4-Oil, Dimilin, and Orthene on Forest Birds in Northeastern Oregon. U.S. Department of Agriculture, Pacific Southwest Forest Range Experiment Station, Research Paper PSW-148. 19 pp.

Sample, B.E., Butler, L., and Whitmore, R.C. 1992. How <u>Bacillus thuringiensis</u> (Bt) Affects Arthropods Other than the Gypsy Moth: Preliminary Results. USDA Forest Service.

Shama, S. K., Etkind, P. H., Odell, T. M., Canada, A. T., Finn, A. M., and Soter, N. A. 1982. Gypsy Moth Caterpillar Dermatitis. <u>The New England Journal of Medicine</u>. 306:21, pp. 1300-01.

Stoner. A., and Wilson, W.T. 1982. Diflubenzuron (dimilin) Effect of Long-Term Feeding of Low Doses in Sugar-Cake or Sucrose Syrup on HoneyBees in Standard-Size Field Colonies. <u>American Bee Journal</u>. 122:579-582.

Tuthill, R.W., Canada, A.T., Wilcock, K., Etkind, P.H., Odell, T. M., and Shama, S.K. 1984. An Epidemiologic Study of Gypsy Moth Rash. <u>American Journal of Public Health</u>. 74:8, pp. 799-803.

- U. S. Department of Agriculture. 1992. Potential Impacts of Asian Gypsy Moth in the Western United States. USDA Forest Service, Forest Pest Management, State and Private Forestry. Washington D.C.
- U. S. Department of Agriculture. 1995. Gypsy Moth Management in the United States: a cooperative approach. Final Environmental Impact Statement as Supplemented--1995. USDA Animal and Plant Health Inspection Service and USDA Forest Service. Washington D.C.

U. S. Environmental Protection Agency. 1998. Reregistration Eligibility Decision (RED) *Bacillus thuringiensis*. EPA738-R-98-004. March 1998, pp. 32-33.


Washington State Department of Health. 1993. Report of Health Surveillance Activities, Asian Gypsy Moth Control Program.

Young, M.F., Trombetta, L.D., and Carson, S. 1986. Effects of Diflubenzuron on the Mouse Liver. <u>Journal of Applied Toxicology</u>. 6:343-348.

APPENDIX B


Treatment Site Maps


Proposed 2004 Gypsy Moth Eradication Area Bellevue (11 Acres)



NOTE: The boundaries for the proposed treatment zone are: on the west, just west of 119th Avenue NE; on the east, just east of 120th Avenue NE: on the north, approximately 425 feet north of NE 8th Street; and on the south, approximately 200 feet south of NE 8th

City of Bellevue MapGuide






SCALE 1: 7,780


500 0 500 1,000 1,500

FEET


UNITED STATES DEPARTMENT OF INTERIOR GEOLOGICAL SURVEY




Proposed 2004 Gypsy Moth Eradication Area Port Ludlow (18 Acres)



NOTE: The boundaries for the proposed treatment zone are: on the west, Osprey Ridge Drive; on the north, Goldfinch Lane; on the east, just west of Rainier Lane; and on the south, just north of Hummingbird Court.







SLIVER LN Mayfield (7.5 Acres) GM Catches 2004 Proposed Treatment Boundaries

Proposed 2004 Gypsy Moth Eradication Area

NOTE: The boundary of the proposed treatment zone is a half-circle with a radius of approximately 450 feet. The half-circle is just west of Flynn Road where 3 pupal cases were found.

Proposed 2004 Gypsy Moth Eradication Area Mayfield (7.5 Acres)





APPENDIX C

Alternatives Described in 1995 FEIS

Alternative 1. No Suppression, No Eradication, No Slow the Spread

Under alternative 1, the Forest Service and APHIS would not suppress, eradicate, or slow the spread of the gypsy moth (fig. 2-5).

Implementation of alternative 1 would not reduce damage, prevent establishment, or slow the spread of the gypsy moth.

Alternative 2. Suppression

Under alternative 2, the Forest Service could conduct suppression projects and cooperate with other Federal agencies and States to conduct suppression projects (fig. 2-6).

The Forest Service and APHIS would not slow the spread in the transition area, and neither would eradicate isolated infestations.

Implementation of alternative 2 would help reduce damage caused by the gypsy moth in the generally infested area.

Alternative 3. Eradication


Under alternative 3 the Forest Service and APHIS could conduct eradication projects and cooperate with other Federal agencies and States to conduct eradication projects (fig. 2-7).


The Forest Service would make no coordinated effort to suppress the gypsy moth in the generally infested area. The Forest Service and APHIS would not slow the spread in the transition area.


Implementation of alternative 3 would prevent establishment of gypsy moth populations in the uninfested area. The Asian strain of the gypsy moth would be eradicated wherever it is found, including the generally infested area when the source of the introduction is known.


Alternative 4. Suppression and Eradication


Under alternative 4 the Forest Service could conduct suppression projects and cooperate with other Federal agencies and States to conduct suppression projects. The Forest Service and APHIS could conduct eradication projects, and cooperate with other Federal agencies and States to conduct eradication projects (fig. 2-8). This alternative proposes the continuation of gypsy moth strategies currently being implemented. Alternative 4 represents the "no action" alternative in that it would be no change from the current program.











USDA agencies would make no coordinated effort to reduce the rate of spread of the insect in the transition area.

Implementation of alternative 4 would reduce damage caused by the gypsy moth in the generally infested area and prevent establishment of gypsy moth populations in the uninfested area. The Asian strain of the gypsy moth would be eradicated wherever it is found, including the generally infested area when the source of the introduction is known.

Alternative 5. Eradication and Slow the Spread

Under alternative 5 the Forest Service and APHIS could conduct eradication and slow-the-spread projects, and cooperate with other Federal agencies and States to conduct eradication and slow-the-spread projects (fig. 2-9).

The Forest Service would make no coordinated effort to suppress outbreak populations of the gypsy moth in the generally infested area.

Implementation of alternative 5 would prevent establishment of gypsy moth populations in the uninfested area and slow the natural spread of the insect in the transition area. The Asian strain of the gypsy moth would be eradicated wherever it is found, including the generally infested area when the source of the introduction is known.

Alternative 6. Suppression, Eradication, and Slow the Spread (Preferred)

Under alternative 6 the Forest Service could conduct suppression projects, and cooperate with other Federal agencies and States to conduct suppression projects. The Forest Service and APHIS could conduct eradication and slow-the-spread projects and cooperate with other Federal agencies and States to conduct eradication and slow-the-spread projects (fig. 2-10). Alternative 6 is the preferred alternative.

Implementation of alternative 6 would help reduce damage in the generally infested area, prevent establishment of gypsy moth populations in the uninfested area, and slow the natural spread of the insect in the transition area. The Asian strain of the gypsy moth would be eradicated wherever it is found, including the generally infested area when the source of the introduction is known.

APPENDIX D

Washington State Department of Health Recommendations



STATE OF WASHINGTON

DEPARTMENT OF HEALTH

OFFICE OF ENVIRONMENTAL HEALTH AND SAFETY

7171 Cleanwater Lane, Building 4 • P.O. 47825 Olympia, Washington 98504-7825 TDD Relay Service (800) 833-6388

February 2004

HEALTH RECOMMENDATIONS FOR GYPSY MOTH SPRAYING

Washington State Department of Agriculture has proposed to use a biological agent *Bacillus thuringiensis var. kurstaki* (Btk) for gypsy moth control in three local areas later this spring. If you live in the area to be sprayed, you are unlikely to experience any health problems due to the spray.

The spray contains bacteria that are natural to soil and have been used for years to kill pest caterpillars. The spray also contains water (99%), a sticking agent that contains synthetic latex (0.12 %), and traces of food crops and preservatives that are approved for use on food. In animal tests, Btk has shown almost no toxicity to mammals, birds, or fish, except at extremely high levels. Many years of experience with Btk products have shown that the vast majority of persons living in sprayed areas have reported no symptoms. Some members of the general public have complained of mild skin reactions; eye, nose, or throat irritation; and worsening of asthma or allergies after aerial spraying.

The Washington Department of Health (DOH) recommends that people in the area to be sprayed minimize exposure by doing the following:

- 1. Stay indoors for at least 30 minutes after the spraying to allow droplets to settle.
- 2. Wait until the spray has dried before letting skin touch the treated leaves and bushes.
- 3. Wash skin with soap and water if you come in contact with the spray.
- 4. People in the sprayed area can sign up with the Department of Agriculture (800-443-6684) to be notified the day before spraying.

Special health concerns: It is possible that individuals with certain health conditions may be more sensitive to the spray. If you have asthma, severe allergies to food or food preservatives, or immune disorders, you may want to stay indoors longer or leave the sprayed area during the day of treatment. We encourage people with special health concerns to talk with their personal doctor for advice.

If you have an illness that you think is related to the spraying, please report this to the Department of Health at (360) 236-3360 or toll free at (888) 586-9427.

For more health information:

Washington State Department of Health
National Pesticide Information Center

Barbara Morrissey
(360) 236-3368
npic@ace.orst.edu
(800) 858-7378



APPENDIX E

Product Labels & Material Safety Data Sheets

Foray 48B

Biological Insecticide Flowable Concentrate

ACTIVE INGREDIENT:

POTENCY: 10,600 International Units (IU)/mg of product (equivalent to 48 billion IU/Gal). Potency units should not be used to adjust use rates.

EPA Reg. No. 73049-46 EPA Est. No. 33762-IA-001

List No. 60178

INDEX:

- 1.0 Statement of Practical Treatment
- 2.0 Precautionary Statements
 - 2.1 Hazard to Humans (and Domestic Animals)
 - 2.2 Personal Protective Equipment (PPE)
 - 2.3 User Safety Recommendations
 - 2.4 Environmental Hazards
- 3.0 Directions for Use
- 4.0 Storage and Disposal
- 5.0 Directions for Non-Agricultural Applications
- 6.0 Mixing
- 7.0 Application
- 8.0 Application Rates
- 9.0 Notice to User

KEEP OUT OF REACH OF CHILDREN CAUTION

For <u>MEDICAL</u> and <u>TRANSPORT</u> Emergencies <u>ONLY</u> Call 24 Hours A Day 1-877-315-9819. For All Other Information Call 1-800-323-9597.

1.0 STATEMENT OF PRACTICAL TREATMENT

If on Skin: Wash with plenty of soap and water. Get medical attention.

If in Eyes: Flush with plenty of water. Call a physician if eye irritation persists.

2.0 PRECAUTIONARY STATEMENTS

2.1 HAZARD TO HUMANS (AND DOMESTIC ANIMALS) CAUTION

Causes moderate eye irritation. Avoid contact with skin, eyes, open wounds or clothing. Wash thoroughly with soap and water after handling.

2.2 Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- * Waterproof gloves
- T Shoes plus socks

Follow the manufacturer's instructions for cleaning/ maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

2.4 Environmental Hazards

Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal of waste.

Storage: Store in a cool, dry place. Keep containers tightly closed when not in use. Store in temperatures above freezing and below 32°C (90°F).

Pesticide Disposal: Pesticide waste resulting from the use of this product may be disposed of on site or at an approved waste disposal facility in accordance with federal and local regulations.

Container Disposal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

5.0 DIRECTIONS FOR NON-AGRICULTURAL APPLICATIONS

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in interior plantscapes, ornamental gardens or parks, or on golf courses or lawns and grounds.

Not for use on trees being grown for sale or other commercial use, or for commercial seed production, or for the production of timber or wood products, or for research purposes except wide-area public pest control programs sponsored by government entities, such as mosquito abatement, gypsy moth control, and Mediterranean fruit fly eradication.

6.0 MIXING

Foray 48B contains the spores and endotoxinal crystals of *Bacillus thuringiensis kurstaki*. Foray 48B is a stomach poison and is effective against lepidopterous larvae. After ingestion, larvae stop feeding within hours and die 2-5 days later. Maximum activity is exhibited against early instar larvae. Foray 48B may be used for both ground and aerial application. The product should be shaken or stirred before use. Add some water to the tank mix, pour the recommended amount of Foray 48B into the tank and then add the remaining amount of water to obtain the proper mix ratio. Agitate as necessary to maintain the suspension. The diluted mix should be used within 72 hours.

7.0 APPLICATION


Ground Application: Use an adequate amount of tank mix to obtain thorough coverage without excessive run off. Use the recommended per acre dosages of Foray 48B in the following amounts of water:

High volume hydraulic sprayers 100 gallons Mist blowers 10 gallons

Aerial Application: Foray 48B may be applied aerially, either alone or diluted with water at the dosages shown in the application rates table. Spray volumes of 32-128 ounces per acre are recommended. Best results are expected when Foray 48B is applied to dry foliage.

APPLICATION RATES

		D-4-4	D
		Rate ¹	Dosage ¹
Crop	Pests	(pts/acre)	(BIU/Acre
Forests,	Gypsy Moth,	1.3-6.7	8-40
Shade Trees,	Asian Gypsy		
Omamentals,	Moth,		
Shrubs,	Elm Spanworm		
Sugar Maple	Spruce Budworm,	1.3-5	8-30
Trees, Ornamental	Browntail Moth,		•
Fruit, Nut	Douglas Fir Tussock Moth,		
and Citrus	Coneworm		
Trees ²	Coneworm		
11000-	Tussock Moths,	1-2.7	6-16
	Pine Butterfly,		
	Bagworm ,		
	Leafrollers,		
	Tortix,		
	.Mimosa		
	Webworm,		
	Tent		
ļ	Caterpillar,		
	Jackpine Budworm,		I
	Blackheaded		
	Budworm,		
	Saddled		
	Prominent,		
	Saddleback		
de de la constante de la const	Caterpillar,		
	Eastern and		
	Western		
	Hemlock		
	Looper,		
	Orange-striped Oakworm.		
	Satin Moth		
	Redhumped	0.7-1.3	4-8
	Caterpillars,		
	Spring and Fall		
	Cankerworm.		
	California		
	Oakworm,		



Fall Webworm

² In treating Gypsy Moth and Asian Gypsy Moth infested trees and shrubs in urban, rural and semi-rural areas, exposure of non-target vegetation including, but not limited to, native and ornamental species and food or feed crops is permitted.

NOTICE OF WARRANTY

9.0

SELLER **MAKES** NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PURPOSE, OTHERWISE. **EXPRESS** OR CONCERNING THIS PRODUCT OR ITS USES WHICH EXTEND BEYOND THE USE OF THE PRODUCT UNDER NORMAL CONDITIONS IN ACCORD WITH THE STATEMENTS MADE ON THIS LABEL. IN NO CASE SHALL THE SELLER BE LIABLE FOR CONSEQUENTIAL, SPECIAL, OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER.

Foray® 48B

1. CHEMICAL PRODUCT AND COMPANY IDENTI-FICATION

MATERIAL NAME: Foray® 48B - Not For Use in Canada

EPA Registration No. 73049-46

List Number: 60175 * 60178 * 60179 * 60180

MANUFACTURER: Valent BioSciences Corporation

870 Technology Way, Suite 100

Libertyville, Illinois 60048

EMERGENCY TELEPHONE NUMBERS

Emergency Health or Spill:

Outside the United States: 1-651-632-6184 Within the United States: 1-877-315-9819

2. COMPOSITION/INFORMATION ON INGREDI-

INGREDIENT NAME: Bacillus thuringiensis, var.

CAS/RTECS NUMBERS: 68038-71-1, N/A

OSHA-PEL BHR TWA: N/L STEL: N/L CEILING: N/L ACGIH-TLV

BHR TWA: N/L

STFL: N/L CEILING: N/L

OTHER LIMITS

8HR TWA: N/A STEL: N/A

CEILING: N/A

INGREDIENT NAME: Inert ingredients - identity with-

held as a Trade Secret

CAS/RTECS NUMBERS: N/A, N/A

OSHA-PEL 8HR TWA: N/L

STEL: N/L CEILING: N/L **ACGIH-TLV**

8HR TWA: N/L

STEL: N/L CFILING: N/L

OTHER LIMITS

8HR TWA: N/A STEL: N/A

CEILING: N/A

EEC (European Community): N/D

Symbol Designation: N/A

Risk Phrases: N/A

Safety Phrases: S2 Keep out of reach of children. S3 Keep in a cool place. S13 Keep away from food, drink and animal feeding stuffs.

3. HAZARDS INFORMATION

EMERGENCY OVERVIEW: This material may cause transient skin and eye irritation.

ROUTE(S) OF ENTRY:

Skin: No Inhalation: No Incestion: No

INGESTION RATING: None SKIN ABSORPTION RATING: None INHALATION RATING: N/D CORROSIVENESS RATING: None SKIN CONTACT RATING: None

SKIN SENSITIZATION RATING: N/D EYE CONTACT RATING: None

TARGET ORGANS: Possibly skin, eyes and respiratory tract

CARCINOGENICITY RATING:

NTP: N/L IARC: N/L OSHA: N/L ACGIH: N/L

SIGNS AND SYMPTOMS: N/D. May cause skin, eye, and respiratory irritation

MEDICAL CONDITIONS AGGRAVATED BY EXPO-SURE: N/D

4. FIRST AID MEASURES

EYES: Remove from source of exposure. Flush with copious amounts of water. If irritation persists or signs of toxicity occur, seek medical attention. Provide symptomatic/supportive care as necessary.

SKIN: Remove from source of exposure. Flush with copious amounts of water. If irritation persists or signs of toxicity occur, seek medical attention. Provide symptomatic/supportive care as necessary.

INGESTION: Remove from source of exposure. If signs of toxicity occur, seek medical attention. Provide symptomatic/supportive care as necessary. INHALATION: Remove from source of exposure. If

signs of toxicity occur, seek medical attention. Provide symptomatic/supportive care as necessary.

5. FIRE FIGHTING PROCEDURES

ELASH POINT: N/D FLASH POINT METHOD: N/D LOWER EXPLOSIVE LIMIT(%): N/D UPPER EXPLOSIVE LIMIT(%): N/D **AUTOIGNITION TEMPERATURE: N/D** FIRE & EXPLOSION HAZARDS: N/D

EXTINGUISHING MEDIA: Use appropriate medium for the underlying cause of the fire

FIRE FIGHTING INSTRUCTIONS: Wear protective clothing and self-contained breathing apparatus.

6. ACCIDENTAL RELEASE MEASURES

SPILL OR RELEASE PROCEDURES: Recover product and place in an appropriate container for disposal. Ventilate and wash the spill area.

7. HANDLING AND STORAGE

HANDLING: Avoid dust generation and provide room ventilation during handling.

STORAGE: Store in a cool, dry place. Keep containers tightly closed when not in use. Store in temperatures above freezing and below 32 C (90 F).

SPECIAL PRECAUTIONS: Wash thoroughly with soap and water after handling.

8. EXPOSURE CONTROLS/PERSONAL PROTEC-TION

ENGINEERING CONTROLS: Use local exhaust RESPIRATORY PROTECTION: Air purifying respirator with dust/mist filter (N95), if necessary. SKIN PROTECTION: Impervious.

EYE PROTECTION: Goggles.

OTHER PROTECTION: Wear tyvek coveralls if contact

9. PHYSICAL AND CHEMICAL PROPERTIES

APPEARANCE/PHYSICAL STATE: Light brown suspension

ODOR: Pungent, bacterial BOILING POINT: N/D MELTING/FREEZING POINT: N/D

VAPOR PRESSURE (mm Hg): N/D VAPOR DENSITY (Air=1): N/D

EVAPORATION RATE: N/D **BULK DENSITY: 1.12-1.2 g/cm3** SPECIFIC GRAVITY: N/D

SOLUBILITY: Readily mixable with water pH: 4.1-4.8 as a 10% solution in water

VISCOSITY: N/D

10. STABILITY AND REACTIVITY

CHEMICAL STABILITY: Not chemically reactive. INCOMPATIBILITIES: Alkalinity inactivates product. HAZARDOUS DECOMPOSITION PRODUCTS: Not

HAZARDOUS POLYMERIZATION: Not known to oc-

11. TOXICOLOGICAL INFORMATION

ORAL TOXICITY: N/D. LD50 > 5000 mg/kg in rats for a similar formulation. EPA Category IV.

DERMAL TOXICITY: N/D. LD50 > 2500 mg/kg in rabbits for a similar formulation. EPA Category III. INHALATION TOXICITY: N/D. A similar formulation was not lethal in an inhalation study at the maximum achievable concentration of 6.81 mg/L. EPA Category CORROSIVENESS: N/D. A similar formulation was not

corrosive.

DERMAL IRRITATION: N/D. A similar formulation had slight skin reactions up to 24 hrs after treatment. EPA Category IV.

OCULAR IRRITATION: N/D. A similar formulation was mildly irritating, which was reversible within 7 days. FPA Category III.

DERMAL SENSITIZATION: N/D SPECIAL TARGET ORGAN EFFECTS: N/D CARCINOGENICITY INFORMATION: N/D

12. ECOLOGICAL INFORMATION

ECOLOGICAL INFORMATION: Studies on nontargets have been performed without identifying any organisms at risk. The following species have been included in the testing. Mammals (rats, rabbits), Freshwater aquatic invertebrate (Daphnia Magna, Rainbow Trout), Birds (Mallard Duck, Bobwhite), Non-target insects (Green Lacewing Larvae, Hymenopteran Pedibus Foveolatus, Ladybird Beetles, Honey Bee).

13. DISPOSAL CONSIDERATIONS

WASTE DISPOSAL METHODS: Dispose of product in accordance with federal, state and local regulations.

14. TRANSPORTATION INFORMATION

DOT

STATUS: Not Regulated PROPER SHIPPING NAME: N/D HAZARD CLASS: N/D UN NUMBER: N/D PACKING GROUP: N/D REPORTABLE QUANTITY: N/D ΙΔΤΔ/ΙζΔΩ

STATUS: Not Regulated

PROPER SHIPPING NAME: N/D HAZARD CLASS: N/D

UN NUMBER: N/D PACKING GROUP: N/D

REPORTABLE QUANTITY: N/D

IMO

STATUS: Not Regulated PROPER SHIPPING NAME: N/D

HAZARD CLASS: N/D UN NUMBER: N/D PACKING GROUP: N/D REPORTABLE QUANTITY: N/D

FLASH POINT: N/D

15. REGULATORY INFORMATION

TSCA STATUS: Exempt CERCLA STATUS: N/D SARA STATUS: N/D RCRA STATUS: N/D PROP 65 (CA): N/D

16. OTHER INFORMATION

LEGEND:

N/A = Not Applicable

N/D = Not Determined N/L = Not Listed

L = Listed C = Ceiling

S = Short-term

(R) = Registered Trademark of Valent BioSciences (TM) = Registered Trademark of Valent BioSciences The information and recommendations contained herein are based upon tests believed to be reliable. However, Valent BioSciences does not guarantee their accuracy or completeness nor shall any of this information constitute a warranty, whether expressed or implied, as to the safety of the goods, the merchantability of the goods, or the fitness of the goods

for a particular purpose. Adjustment to conform with actual conditions of usage may be required. Valent BioSciences assumes no responsibility for results obtained or for incidental or consequential damages arising from the use of these data. No freedom from infringement of any patent, copyright or trademark is

to be inferred. © Valent BioSciences Corporation June 2001





CAUTION: Keep out of Reach of Children NET CONTENTS: 1 U.S. Gallon (3.785 Litres)

CAUTION: Harmful if absorbed through skin, Causes moderate eye irritation. Avoid contact with skin, eyes or clothing, Wash thoroughly with soap and water after handling. Personal Protective Equipment: Wear Long-sleeved shirt and long pants, Socks, Shoes and

First Aid: If on skin of clothing:

Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes

Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present,

If in eyes: after the first 5 minutes, then continue rinsing eye.

Call a poison control center or doctor immediately for treatment advice. Have a person sip a glass of water if If swallowed:

able to swallow. Do not give anything by mouth to an unconscious person.

Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artifical respiration, If inhaled: preferably by moth-to-mouth, if possible.

General: BOND is a very efficient sticker for agriculture and can be used in terrestrial or aquatic settings. BOND's adhesion properties increase initial deposition, reduces run-off and secures spray from rain or overhead irrigation. Apply sprays containing BOND at least one hour before an anticipated rain or overhead irrigation. Once the spray has dried, BOND will adhere the pesticides.

Directions for Use: SHAKE WELL BEFORE USE. Fill spray tank 1/2 full with water and begin agitation. Add pesticides as directed by the label while maintaining agitation and continue to fill. After pesticides are thoroughly mixed, continue agitation and add BOND at desired rate. Some posticides have stated adjuvant use rates. In all cases, the pesticide manufacturer's label should be consulted regarding specific use recommendations and that rate followed. Do not add adjuvant at a level that would exceed 5% of the finished

Suggested use rates: The sticking efficiency of BOND varies from pesticide to pesticide, so the usage rate will be associated with the formulation being sprayed.

1 to 2 pints per 100 gallons OR 2 to 4 fluid ounces per acre

Rinse tank and nozzles immediately after spraying. Observe the pre-harvest interval on the pesticide label when using BOND. No time limitations apply to non-food crops.

Storage: Store in cool, dry place. Store in original container. Keep tightly closed. Do not reuse empty container. Disposal: Do not contaminate water, food or feed by storage or disposal. Wastes may be disposed of on-site or at an approved waste disposal facility. Triple rinse (or equivalent) adding rinse water to spray tank. Offer container for recycling or dispose of container in sanitary landfill, or by other procedures approved by appropriate authorities. Recycling decontaminated containers is the best option of container disposal. The Agricultural Container Recycling Council (ACRC) operates the national recycling program. To contact your state and local ACRC recycler visit the ACRC web page at www.acrecycle.org.

WARRANTY DISCLAIMERS AND NOTICE
IT IS IMPOSSIBLE TO ELIMINATE ALL RISKS INHERENTLY ASSOCIATED WITH THE USE OF THIS PRODUCT. CROP INJURY, INEFFECTIVENESS, OR OTHER UNINTENDED CONSEQUENCES MAY RESULT BECAUSE OF SUCH FACTORS AS WEATHER CONDITIONS, PRESENCE OF OTHER MATERIALS, OR THE MANNER OF USE OR APPLICATION, ALL OF WHICH ARE BEYOND THE CONTROL OF LOVELAND INDUSTRIES, INC., THE MANUFACTURER, OR SELLER. IN NO CASE SHALL LOVELAND INDUSTRIES, INC., THE MANUFACTURER, OR SELLER BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER OR USER.

RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. ALL SUCH HISKS STALL BE ASSUMED BY THE BUTCH OF USER.

EXCEPT AS EXPRESSLY PROVIDED HEREIN, LOVELAND INDUSTRIES, INC., THE MANUFACTURER, OR SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO, MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. BUYER'S OR USER'S EXCLUSIVE REMEDY, AND LOVELAND INDUSTRIES, INC.'S THE MANUFACTURER'S OR SELLER'S TOTAL LIABILITY, SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT. NO AGENT OR EMPLOYEE OF LOVELAND INDUSTRIES, INC. OR SELLER SHALL BE ABLE TO AMEND THE TERMS OF THIS LABEL.

а

.

BOND

I. IDENTIFICATION OF PRODUCT

LOVELAND INDUSTRIES, INC. P.O. BOX 1289

GREELEY, COLORADO 80632

EMERGENCY PHONE NUMBERS 1-800-424-9300 CHEMTREC

1-970-356-8920 LOVELAND IND.

CALL CHEMTREC 24 HOURS A DAY @ 1-800-424-9300

TRADE NAME: BOND

CHEMICAL NAME: Carboxylated Synthetic Latex (combination synthetic latex and primary aliphatic oxyalkylated alcohol)

II. HAZARDOUS COMPONENTS OF MIXTURES

COMPONENT NONE CAS#

%

TLV(UNITS)

III. PHYSICAL DATA

BOILING POINT: 212'F
BULK DENSITY: 8.41#/Gal
SPECIFIC GRAVITY: 1.01 g/ml
SOLUBLE IN WATER: Yes
VAPOR PRESSURE (mm OF Hg) @ 20 D C: Not Est.
VAPOR DENSITY (AIR=1): Not Est.
% VOLATILE BY VOL: 45%
APPEARANCE: White Emulsion
ODOR: Slight Ammonia
pH: (neat) 7 - 8

IV. FIRE & EXPLOSION HAZARD DATA

FLASH POINT (TEST METHOD): Not Est. FLAMMABLE LIMITS (UEL AND LEL): Not Est. EXTINGUISHING MEDIA: Water spray/fog, Foam, Dry Chemical, CO₂.

SPECIAL FIRE FIGHTING PROCEDURES: Wear protective clothing.

UNUSUAL FIRE AND EXPLOSION HAZARD: None

V. REACTIVITY DATA

STABILITY: Stable

CONDITIONS TO AVOID: None

INCOMPATIBILITY (MATERIALS TO AVOID): Avoid multivalent transition metal ions as they will cause coagulation. Will also coagulate under low pH conditions. HAZARDOUS DECOMPOSITION PRODUCTS: None Known

HAZARDOUS POLYMERIZATION: Will not occur

VI. HEALTH HAZARD DATA

EFFECTS OF OVEREXPOSURE: Animal studies indicate minimal toxicity through oral or dermal route. Eye and skin exposure indicated as minimally irritating. EMERGENCY AND FIRST AID PROCEDURES:

EYES: Flush with water for 15 minutes, then get medical attention.

SKIN: Remove contaminated clothing. Wash with soap and water. Get medical attention if irritation persists.

INGESTION: First aid is not normally required, if symptoms persist get medical attention.

INHALATION: Remove victim to fresh air. Apply artificial respiration if necessary.

CARCINOGEN STATUS: Not listed by NTP, IARC or ACGIH.

VII. SPILL OR LEAK PROCEDURES

STEPS TO BE TAKEN IF MATERIAL IS RELEASED OR SPILLED: Pick up with absorbent material and place in a container for proper disposal in accordance with all Federal, Slate and Local Regulations.

WASTE DISPOSAL METHOD: Do not contaminate water, food or feed by storage or disposal. Dispose of in an approved waste disposal facility in accordance with all Federal, State and Local Regulations.

CONTAINER DISPOSAL: Triple rinse (or equivalent) adding rinse water to spray tank. Offer container for recycling or dispose of in a sanitary landfill or by other procedures approved by the appropriate authorities.

VIII. SPECIAL PROTECTION INFORMATION

RESPIRATORY PROTECTION: Wear NIOSH/MSHA approved respirator if necessary. VENTILATION:

LOCAL: Recommended
MECHANICAL: Not required
SPECIAL: None

OTHER: None

PROTECTIVE GLOVES: Wear rubber or impervious oloves.

gloves.
EYE PROTECTION: Wear goggles or a face shield.
OTHER PROTECTION: Full body covering clothing.

IX. SPECIAL PRECAUTIONS

PRECAUTIONS TO BE TAKEN IN STORAGE AND HANDLING: Store in a cool dry place. Keep in original container, tightly closed. Do not reuse empty container

OTHER PRECAUTIONS: Keep out of reach of children. Always launder contaminated clothing before reuse.

X. REGULATORY INFORMATION

Components which could require reporting under SARA TITLE III are: None.
SARA TITLE III HAZARD CATEGORY

IMMEDIATE: Yes

FREIGHT CLASS: ITEM 102120 CLASS 60

DOT REGULATION: Not Regulated

FIRE: No

REACTIVITY: No

SUDDEN RELEASE OF PRESSURE: No

Legal responsibility is assumed only for the fact that all studies reported here and all opinions are those of qualified experts. Buyer assumes all risks & liability. He accepts & uses this material on these conditions. He must have a copy of this MSDS where this material is handled.

Date of Issue: 3-13-96

Supersedes: 10-25-90

APPENDIX F Standard Operating Procedures

WASHINGTON STATE DEPARTMENT OF AGRICULTURE

STANDARD OPERATING PROCEDURES

2004 Gypsy Moth Eradication Project

- 1. The health and safety of the public, employees of the contractor, and employees of the Washington State Department of Agriculture will be the first concern in implementing the project.
- 2. Mixing and application of the insecticide will be done only by an appropriately licensed applicator and will be done only under the supervision of a Washington State Department of Agriculture treatment site monitor.
- 3. The insecticide will be applied according to label directions.
- 4. Residents in the affected eradication area will be notified of the projected dates and times of insecticide applications through direct mailings, open house presentations, and press releases. Additionally, a manned 1-800 hotline will be established to address further resident concerns, comments, and project suggestions. Recommendations concerning health and welfare issues will be included in public outreach efforts.
- 5. The project will commence at the appropriate stage of leaf and/or larval development.
- 6. Weather conditions, particularly wind, will play the largest role in determining when an effective treatment can be made. In the event of rainfall before spray has had sufficient time to adhere to the foliage, a re-treatment may be necessary.
- 7. Spill control kits will be on site and readily available during all applications.
- 8. Treatments will not occur when wind speed exceeds 10 miles/hour.
- 9. Application over surface waters not associated with trees and shrubs will be avoided.
- 10. Hydraulic apparatus pressures will be limited to that necessary to obtain thorough coverage to the tops of the tallest trees within the treatment area.

APPENDIX G

Letters and Permits Received Through Interagency Consultation Concerning Threatened, Endangered, and Sensitive Species and Habitats

January 15, 2004

Chad Phillips
Plant Protection Division – Entomology Branch
Department of Agriculture
3939 Cleveland Ave SE
Olympia WA 98501

SUBJECT: Three Gypsy Moth Eradication Projects: Bellevue (T25N R05E S28,33); Port Ludlow (T28N R01E S17); and Mayfield (T13N R02E S35)

We've searched the Natural Heritage Information System for information on significant natural features in your project areas. Currently, we have no records for rare plants or high quality native ecosystems in the vicinity of your projects.

The information provided by the Washington Natural Heritage Program is based solely on existing information in the database. In the absence of field inventories, we cannot state whether or not a given site contains high quality ecosystems or rare plant species; there may be significant natural features in your study areas of which we are not aware.

The Washington Natural Heritage Program is responsible for information on the state's rare plants as well as high quality ecosystems. For information on animal species of concern, please contact Priority Habitats and Species, Washington Department of Fish and Wildlife, 600 Capitol Way N, Olympia WA 98501-1091, or by phone (360) 902-2543.

Please visit our internet website at http://www.dnr.wa.gov/nhp for more information. Lists of rare plants and their status, as well as rare plant fact sheets, are available for download from the site. Please feel free to call me at (360) 902-1667 if you have any questions, or by e-mail at sandra.moody@wadnr.gov.

Sincerely,

Sandy Swope Moody, Environmental Review Coordinator

Washington Natural Heritage Program

Sandy Suipe Moody

Asset Management & Protection Division, PO Box 47014, Olympia WA 98504-7014 FAX 360-902-1789



State of Washington DEPARTMENT OF FISH AND WILDLIFE

Mailing Address: 600 Capitol Way N • Olympia, WA 98501-1091 • (360) 902-2200, TDD (360) 902-2207 Main Office Location: Natural Resources Building • 1111 Washington Street SE • Olympia, WA

February 12, 2004

Mr. Chad Phillips Washington State Department of Agriculture Post Office Box 42560 Olympia, WA 98504-2560

Dear Mr. Phillips:

This letter is in response to your December 12, 2003 request to review our butterfly records for the Washington State Department of Agriculture (WSDA) proposed 2004 gypsy moth (*Lymantria dispar*) eradication. We have reviewed our butterfly data and evaluated site habitat conditions for the proposed Bellevue, Port Ludlow, and Mayfield treatment areas. According to your request letter, the proposed Bellevue treatment area is approximately 11 acres and located in T25N R5E S28 and S33; Port Ludlow is approximately 18 acres and in T28N R1E S17; Mayfield is located in T13N R2E S35, and according to our December 30, 2003 telephone discussion, is now approximately 7.5 acres in size. We understand that evidence of gypsy moth reproduction (egg cases) has been found at both the Bellevue and Port Ludlow sites, and pupal cases have been located at the Mayfield site. According to your letter, WSDA proposes as many as four ground-based applications of *Bacillus thuringiensis* var. *kurstaki* (Btk) at each of the sites.

We have reviewed our butterfly records and evaluated local landscape and site habitat conditions at each of the treatment areas for their potential to support rare, state candidate or state listed butterflies. We found no butterfly species of concern records in the immediate proposed Btk application areas or within a 5-mile radius of the areas. Habitat conditions were assessed using aerial photos and WSDA site descriptions. The vegetation conditions at and near these sites make it highly unlikely that they could support rare lepidopterans. The Bellevue proposed treatment site is highly urbanized, the Port Ludlow site is in a high-density housing development, and the Mayfield site is located in a dairy farm's residence yard and barnyard.

We are generally cautious about the use of Btk, due to the potential for impacting local non-target lepidopterans, particularly low-dispersing species that are isolated or patchily distributed. However, given the habitat conditions present at the proposed treatment sites, it is unlikely that such species inhabit these areas. Direct effects on non-target lepidopterans and any associated indirect effects on non-target vertebrates are likely to be minimal and short-term as the application areas are small and habitat within the areas is similar to the surrounding landscape, factors that support lepidopteran recolonization. We recognize the importance and support early

Mr. Chad Phillips February 12, 2004 Page 2

eradication of gypsy moth when populations become established in Washington. We encourage WSDA's participation in ongoing research to develop effective gypsy moth treatment methods that are less harmful to non-target Lepidoptera.

I hope this information is helpful. If you have any further questions, please contact me at 360-902-2496.

Sincerely,

Am E. Potter

Ann E. Potter, Wildlife Biologist

Wildlife Diversity Division

AEP:aep:ahr

cc: Fred Dobler

Carl Dugger

Jack Smith

Steve Kalinowski

Lora Leschner

Rich Costello

Robert M. Pyle

Richard Youel